arXiv:1302.4004v1 [math-ph] 16 Feb 2013

GENERALIZING THE CONNES MOSCOVICI HOPF ALGEBRA TO CONTAIN ALL ROOTED TREES

SUSAMA AGARWALA AND COLLEEN DELANEY

Abstract. This paper defines a generalization of the Connes-Moscovici Hopf algebra, $\mathcal{H}(1)$ that contains the entire Hopf algebra of rooted trees. A relationship between the former, a much studied object in non-commutative geometry, and the later, a much studied object in perturbative Quantum Field Theory, has been established by Connes and Kreimer. The results of this paper open the door to study the cohomology of the Hopf algebra of rooted trees.

The Connes-Moscovici defina a family of Hopf algebras, $\mathcal{H}(n)$, on n-dimensional flat manifolds in [6]. The action of these Hopf algebras on the group of smooth functions on the frame bundle crossed with local diffeomorphism on M, is a well studied object in non-commutative geometry,

In [5], Connes and Kreimer delineate a remarkable set of similarities between the Hopf algebra of rooted trees, $\mathcal{H}_{rt}$, which is important to understanding the perturbative regularization of Quantum Field Theories (QFTs), and $\mathcal{H}(1)$. The Hopf algebra $\mathcal{H}(1)$ can be defined by a Lie algebra generated by two vector fields on the orientation preserving frame bundle F^+M, X and Y, and a set of linear operators $\{\delta_i | i \in \mathbb{N}\}$. The linear operators correspond to a sub Hopf algebra of $\mathcal{H}_{rt}$, the vector field Y corresponds to the grading operator on $\mathcal{H}_{rt}$ and the vector field X to the natural growth operator on $\mathcal{H}_{rt}$.

While a lot is known about the cohomologies of $\mathcal{H}(n)$, little is known about the cohomologies of $\mathcal{H}_{rt}$. However, one of the few cocycles of the Hochschild cohomology that is well known, the grafting operator B_+, plays an important role in understanding the combinatorics underlying the Dyson Schwinger equations for perturbative QFTs [1, 8, 9]. Knowing more about cohomology of $\mathcal{H}_{rt}$ will lead to a better understanding of the process of perturbative regularization of QFTs, as well as the deep connection between non-commutative geometry and renormalization of the same. Framing the full Hopf algebra of rooted trees in the context of $\mathcal{H}(1)$, is a first step in this direction.

This paper introduces a generalization of $\mathcal{H}(1)$, $\mathcal{H}_{rt}(1)$, that contains all of $\mathcal{H}_{rt}$ as a sub Hopf algebra. Key to this construction is an observation by Cayley in 1881 relating rooted trees to a particular first order differential equation. We use this to generalize the natural growth operator on $\mathcal{H}_{rt}$, and thus to define a family of vector fields X_t on F^+M, corresponding to the generalized natural growth operators. The commutation of these vector fields X_t with the element δ_1 in $\mathcal{H}(1)$ gives a family of linear operators δ_t, each corresponding to a rooted tree $t \in \mathcal{H}_{rt}$. In order to build this Hopf algebra $\mathcal{H}_{rt}(1)$, we have to relax the requirement that the manifold M is flat.

This paper is organized as follows. Section 1 recalls the construction of Cayley, and the relationship between differentials and rooted trees. Section 2 recalls the definition of $\mathcal{H}_{rt}$, and the natural growth operator. We generalize the natural growth operator, and show that all of $\mathcal{H}_{rt}$ can be generated by repeated application of the generalized natural growth operator to the single vertex tree. Section 3 recalls the definition of $\mathcal{H}(1)$ and constructs the generalized Hopf algebra $\mathcal{H}_{rt}(1)$.

Acknowledgments

The authors would like to thank Karen Yeats for several useful discussions during the early stages of this project. We are also grateful to the Caltech SURF program for partial funding for this project.

1. A relationship between rooted trees and differentials

A relationship between rooted trees and differentials has been known since Cayley's work in 1881 [4]. This relationship is fundamental to Butcher's work to to solve differential equations using the Runge Kutta method [3]. For a brief but clear exposition of the results in this section, see [2]. Let $\mathbf{x}(s)$ be a vector field and

$$\frac{d}{ds}\mathbf{x}(s) = \mathbf{f}(\mathbf{x}) \qquad \mathbf{x}(0) = \mathbf{x}_0 \ . \tag{1}$$

the system of differential equation of interest to Butcher. The Butcher group for this system of differential equations is the Lie group defining the Hopf algebra of rooted trees. Specifically, let $\mathcal{H}_{rt}$ be the Hopf algebra of rooted trees, defined in section 2 of this paper. Let G be the Butcher group for this differential equation. Then $\mathcal{H}_{rt}$ is the graded dual of the universal enveloping algebra of the Lie algebra of G

$$\mathcal{H}_{rt} = \mathcal{U}^\vee(\mathrm{Lie}(G)) \ .$$

The group G defines the combinatorics at the heart of $\mathcal{H}_{rt}$. For more details, see [7]. For a detailed exposition on the relationship between the Butcher group and the Hopf algebra of rooted trees, see [2].

For the rest of this paper, we consider differential equations of the form (1). For now, let $\mathbf{x}(s)$ be a smooth vector field

$$\mathbf{x}(s) : \mathbb{R} \to \mathbb{R}^n \ .$$

In section 3, we will generalize these arguments from $\mathbb{R}^n$ to more general manifolds.

Definition 1. Write $\partial_i = \frac{\partial}{\partial x^i}$. We use the shorthand $\partial_{i_1 \ldots i_k} f(\mathbf{x}) := \partial_{i_1}\partial_{i_2}\ldots\partial_{i_k} f(\mathbf{x})$.

In the context where $\mathbf{x}(s)$ is a vector field on $\mathbb{R}^n$, the following correspondence exists between derivations of $\mathbf{x}(s)$ and non-planar rooted trees.

Definition 2. Rooted trees are non-planar simply connected graphs with a marked point, called the root. The edges of the graph are oriented away from the root.

In this case, rooted trees are drawn root vertex up, indicated by a circle around the root vertex.

$\frac{d}{ds}\mathbf{x}(s) = \mathbf{f}(\mathbf{x})$



$\frac{d^2}{ds^2}\mathbf{x}(s) = \frac{d}{ds}\mathbf{f}(\mathbf{x}) = D\mathbf{f}(\mathbf{x}) \cdot \frac{d}{ds}\mathbf{x}(t) = \partial_i f^j(\mathbf{x}) \frac{d}{ds} x^i(s)$



$\frac{d^3}{ds^3}\mathbf{x}(s) = \frac{d}{ds}(\partial_i f^j(\mathbf{x}))f^i + \partial_i f^j \frac{d}{ds} f^i = (\partial_{ik} f^j) f^k f^i + (\partial_i f^j)(\partial_k f^i) f^k$



$\frac{d^4}{ds^4}\mathbf{x}(s) = (\partial_{ikl} f^j) f^l f^k f^i + 2(\partial_{ik} f^j)(\partial_l f^i) f^l f^k + (\partial_{li} f^j)(\partial_k f^i) f^l f^k + (\partial_i f^j)(\partial_{lk} f^i) f^l f^k + (\partial_i f^j)(\partial_k f^i)(\partial_l f^k) f^l$



Notice that the n^{th} derivative of $\mathbf{x}(s)$ with respect to s gives rise to a linear combination of trees witn n vertices. In fact, it is a linear combination of all the trees with n vertices.

Definition 3. Let $V(t)$ be the vertex set of a tree, and $E(t)$ the set of edges. If v and $w \in V(t)$ are the initial and terminal vertices, respectively, of an edge in t, indicate this edge as $vw \in E(t)$. The *fertility* of a vertex, v, is the number of edges for which v is the initial vertex. Write this $fert(v)$. If $fert(v) = 0$, then v is a leaf vertex.

Notice that each vertex $w \in V(t)$ is either the root vertex, or there is a unique vertex $v_w \in V(t)$ such that $v_w w \in E(t)$. The vertex v_w is called the parent vertex of w. For a fixed differential equation as in (1) above, following Butcher [3], one can associate to any rooted tree t a function of f and its derivatives,

$$\phi : \mathcal{H}_{rt} \to C^\infty(\mathbb{R}^n, \mathbb{R}^n) \ . \tag{2}$$

To fix notation, define $\frac{d}{ds}x^i = f^i = \phi^i(\bullet)$. The map ϕ is defined on rooted trees, where r is the root vertex of t,

$$\phi^i(t) = \Big(\prod_{w\in V(t)\backslash r} \partial_{w_1\dots w_{fert(w)}}\phi^{v_{w_k}}(\bullet)\Big)\partial_{r_1\dots r_{fert(r)}}\phi(\bullet)^i \ . \tag{3}$$

For instance, given a tree

$t =$



one has

$$\phi^k(t) = \phi^j_\bullet\phi^l_\bullet(\partial_l\phi^i(\bullet))(\partial_{ij}\phi^k(\bullet)) \ .$$

It will be useful later in this paper to define a family of linear maps on $C^\infty(\mathbb{R}^n)$ indexed by rooted trees.

Definition 4. Define a linear map

$$\begin{aligned}\phi_t : C^\infty(\mathbb{R}^n) &\to C^\infty(\mathbb{R}^n)\\ h &\to \Big(\prod_{w\in V(t)\backslash r}\partial_{w_1\dots w_{fert(w)}}\phi(\bullet)^{v_{w_k}}\Big)\partial_{r_1\dots r_{fert(r)}}h \ .\end{aligned}$$

In this notation, $\phi(t)^i = \phi_t(\phi(\bullet)^i)$ and $\phi_\bullet(h) = h$.

2. The Hopf algebra of rooted trees

Let $\mathcal{H}_{rt}$ be the Hopf algebra of rooted trees. This Hopf algebra is defined in detail in [5] and summarized here. Consider the vector field generated by all rooted trees t

$$\mathbb{Q}\langle t | t \text{ rooted tree}\rangle \ .$$

This is a unital algebra with the unit representing the empty tree, $1 = \mathbb{I}_{\mathcal{H}_{rt}} = t_\emptyset$. Multiplication on this algebra is commutative, and corresponds to the disjoint union of trees. This is called a forest.

The algebra $\mathcal{H}_{rt}$ is graded by the number of vertices in the tree

$$\mathcal{H}_{rt} = \bigoplus_{i=0}^{\infty}\mathcal{H}_{rt,i}; \quad \mathcal{H}_{rt,0} = \mathbb{Q}; \quad \mathcal{H}_{rt,i} = \mathbb{Q}\langle\{t | t \text{ tree or forest with } i \text{ vertices}\}\rangle \ .$$

There is a grading operator Y on $\mathcal{H}_{rt}$ such that for a single tree with $|V(t)| = i$ vertices, $t \in \mathcal{H}_{rt,i}$, $Y(t) = it$. Before defining the coproduct structure on $\mathcal{H}_{rt}$, we define a few terms.

Definition 5. A proper admissible cut of a tree t is a collection of edges, $c \in E(t)$, such that any path from a root vertex to a leaf vertex contains at most one such edge.

Removing the edges corresponding to an admissible cut of t gives a set of subtrees of t.

Definition 6. The subtree of t associated to a proper admissible cut c that contains the root vertex of t is the root subtree defined by c, $R_c(t)$. The other set of subtrees is the pruned forest defined by$c$, $P_c(t) = \prod_i P_{c,i}(t)$. Each $P_{c,i}(t)$ is a single tree in this subset. The root vertex of each $P_{c,i}(t)$ is defined by the orientation of t.

The set of admissible cuts of a tree consists of the proper admissible cuts and two trivial cuts, one such that $R_c(t) = t$ (called the *empty cut*), and one such that $P_c(t) = t$ (called the *full cut*).

The bialgebra structure on $\mathcal{H}_{rt}$ is given by a coproduct

$$\Delta(t) = \sum_{c \text{ admis. cut}} P_c(t) \otimes R_c(t) \ .$$

This coproduct is compatible with multiplication on $\mathcal{H}_{rt}$. Let c be an admissible cut of t and c' an admissible cut of t'. Then

$$\Delta(tt') = \sum_{c \text{ admis. cut}} \sum_{c' \text{ admis. cut}} P_c(t)P_{c'}(t') \otimes R_c(t)R_{c'}(t) \ .$$

This is a grading preserving coassociative coproduct. The counit is given by

$$\varepsilon(t) = \begin{cases} t & t \in \mathcal{H}_{rt,0}; \\ 0 & \text{else.} \end{cases}$$

Thus defined, $\mathcal{H}_{rt}$ is a connected graded bialgebra over a commutative ring, $\mathbb{Q}$. Therefore, it is a Hopf algebra. The antipode is given by

$$S(t) = -t - \sum_{c \text{ proper admis. cut}} P_c(t)S(R_c(t)) \ .$$

The map ϕ defined in (2) is an algebra homomorphism. For a forest tt',

$$\phi^i(tt') = \phi^i(t)\phi^i(t') \ .$$

Furthermore

$$\Delta\phi^i(t) = \phi^i \otimes \phi^i \Delta t \ .$$

2.1. **The genralized natural growth operator.** In [5] the authors define a sub Hopf algebra of $\mathcal{H}_{rt}$ generated by elements formed by repeated application of the natural growth operator on the single vertex tree. The construction of this Hopf algebra is summarized here, and the idea of natural growth is generalized.

Definition 7. The natural growth operator

$$N : \mathcal{H}_{rt,i} \to \mathcal{H}_{rt,i+1}$$

maps a tree, t, to a sum of trees, $N(t)$, formed by increasing the fertility of each vertex of t by one. Each tree in the sum has one more vertex than t.

For instance,



In loc. cit., the authors define a set of generators

$$N^{k-1}(\bullet) = \delta_k,$$

and check that the algebra $\mathbb{Q}[\{\delta_k | k \in \mathbb{N}\}]$ is a Hopf algebra.

Below are the trees for the first few δ_k.

$\delta_1 =$

$\delta_2 =$ 

$\delta_3 =$ 

$\delta_4 =$  $+\, 3$  $+$  $+$ 

Definition 8. Let $\mathcal{H}_{CK} = \mathbb{Q}[\{\delta_k | k \in \mathbb{N}\}]$ be the Hopf algebra defined in [5], generated by the terms δ_k.

In the context of section 1, given an initial value problem of the form (1), the natural growth operator corresponds to derivation by the parameter s. That is,

$$\phi(\delta_k)^i = \frac{d^k}{ds^k} x^i(s) \ .$$

Lemma 1. *In general, one has the relation*

$$\phi^i(N(t)) = \frac{d}{ds}\phi^i(t) \ .$$

Proof. Recall that for a fixed differential equation

$$\frac{d}{ds}\mathbf{x}(s) = \mathbf{f}(\mathbf{x}) \ ,$$

and a tree $t \in \mathcal{H}_{rt}$, we have

$$\phi(t)^i = \partial_{r_1 \ldots r_{fert(r)}} \phi(\bullet)^i \prod_{w \in V(t) \setminus r} \partial_{w_1 \ldots w_{fert(w)}} \phi(\bullet)^{v_{w_k}} \ .$$

By definition of the natural growth operator,

$$\phi^i(N(t)) = (\partial_{r_1 \ldots r_{fert(r)+1}} \phi(\bullet)^i) \phi(\bullet)^{r_{fert(r)+1}} (\prod_{w \in V(t) \setminus r} \partial_{w_1 \ldots w_{fert(w)}} \phi(\bullet)^{v_{w_k}}) +$$
$$\sum_{u \in V(t) \setminus r} (\partial_{r_1 \ldots r_{fert(r)}} \phi(\bullet)^i)(\partial_{u_1 \ldots u_{fert(v)+1}} \phi(\bullet)^{v_{u_k}}) \phi(\bullet)^{u_{fert(u)+1}} \prod_{w \in V(t) \setminus \{r,u\}} \partial_{w_1 \ldots w_{fert(w)}} \phi(\bullet)^{v_{w_k}} \ .$$

This is exactly the expression for $\frac{d}{ds}\phi^i(t)$. ☐

On the level of trees, there is no reason to limit natural growth only to growth by one vertex. One can grow a tree by any other tree.

Definition 9. For any tree $t \in \mathcal{H}_{rt}$, there is a generalized natural growth operator

$$N_t : \mathcal{H}_{rt,i} \to \mathcal{H}_{rt,i+Y(t)}$$

that maps a tree to a sum of trees, each formed by by increasing the fertility of a vertex by one. This additional edge connects the vertex v of the original tree to the root vertex of t. We define

$$N_t(1) = 0 \text{ and } N_1(t) = Y(t) \ .$$

For instance, let

$t =$  .

Then the tree

$N_t(\delta_2) =$



$+$



.

Lemma 2. *In general, for t and $t' \in \mathcal{R}$,*

$$\phi^i(N_t(t')) = \phi^j(t)\partial_j(\phi^i(t')) \ . \tag{4}$$

Proof. This proof is a generalization of the statement for $t = \delta_1$. Natural growth by t increases the fertility of each vertex by one. This is represented in the right hand side of (4) by differentiation with respect to $x^j(s)$. Instead of growing by a single vertex, N_t grows by the tree t. This is represented by contraction with $\phi^j(t)$. □

The natural growth operator can be extended as an algebra homomorphism $N_t : \mathcal{H}_{rt} \to \mathcal{H}_{rt}$. Specifically

$$N_t(t_1 t_2) = N_t(t_1)N_t(t_2) \text{ and } N_t(t_1 + t_2) = N_t(t_1) + N_t(t_2) \ .$$

Finally, I calculate the composition of the coproduct with the generalized natural growth operator.

One can generalize natural growth to an operator on $C^\infty(\mathbb{R})$:

$$\phi_{N_t} = \phi^j(t)\partial_j \ . \tag{5}$$

Theorem 1. *Let $t, s \in \mathcal{R}$. The coproduct of the operator N_t is given by*

$$\Delta \circ N_t(s) = (N_t \otimes \mathbb{I}) \circ \Delta s + \sum_{c \ admis. \ cut} (P_c(t) \otimes N_{R_c(t)})\Delta s \ .$$

Proof. Let s_v be the tree built by connecting the tree s to t by an edge from the vertex $v \in V(s)$ to the root vertex $r \in V(t)$. Then the set $E(s_v) = E(t) \cup E(s) \cup vr$, and $N_t(s) = \sum_v s_v$. Therefore,

$$\Delta N_t(s) = \sum_v \Delta s_v \ .$$

The set of admissible cuts of s_v can be divided into two sets,

$$A = \{c \text{ admis. cut } | c \cap (E(t) \cup vr) = \emptyset\} \text{ and } B_v = \{c \text{ admis. cut } | c \cap (E(t) \cup vr) \neq \emptyset\} \ .$$

The set A is just the set of admissible cuts of s.

In this notation,

$$\Delta N_t(s) = \sum_{c \in A} \sum_v P_c(s_v) \otimes R_c(s_v) + \sum_v \sum_{c \in B_v} P_c(s_v) \otimes R_c(s_v) \ .$$

For $c \in A$, if the vertex v lies in a pruned subtree, $P_{c,1}(s_v)$, write

$$P_{c,i}(s_v) = (P_{c,i}(s))_v \ .$$

If the vertex v lies in $R_c(s_v)$, write

$$R_c(s_v) = (R_c(s))_v \ .$$

By construction, the terms

$$N_t(P_{c,i}(s_v)) = \sum_{v \in V(P_{c,i}(s_v))} (P_{c,i}(s))_v \quad \text{and} \quad N_t(R_c(s_v)) = \sum_{v \in V(R_c(s_v))} (R_c(s))_v \ .$$

After rearranging terms,

$$\sum_{c \in A} \sum_v P_c(s_v) \otimes R_c(s_v) = (N_t \otimes \mathbb{I}) \circ \Delta(s) + (\mathbb{I} \otimes N_t) \circ \Delta(s) \ . \tag{6}$$

For each $c \in B_v$, write $c = c_s \cup c_t$ with c_t corresponding to an admissible cut of t and c_s an admissible cut of s. If $vr \in c$, then consider $vr \in c_t$ as the empty cut of t. For a fixed $c \in B_v$,

$$P_c(s_v) \otimes R_c(s_v) = P_{c_s}(s)P_{c_t}(t) \otimes R_c(s_v) \ .$$

The root subtree of each $c \in B_v$, $R_c(s_v)$ is formed by attaching the tree $R_{c_t}(t)$ to $R_{c_s}(s)$ by an edge from v to the root vertex of $R_{c_t}(t)$. It is a summand of $N_{R_{c_t}(t)}R_{c_s}(s)$. Notice that by construction of B_v, if $c \in B_v$ it cannot contain the empty cut of s. Again, by rearranging terms

$$\sum_v \sum_{c \in B_v} P_c(s_v) \otimes R_c(s_v) = \sum_{c_s \neq \text{empty}} P_{c_s}(s)P_{c_t}(t) \otimes N_{R_{c_s}(s)}(t) \ . \tag{7}$$

Combining (6) and (7), gives

$$(N_t \otimes \mathbb{I}) \circ \Delta(s) + \sum_{c \text{ admis. cut of } t} (P_c(t) \otimes N_{R_c(t)}) \circ \Delta(s)$$

as desired. $\square$

2.2. **Generating $\mathcal{H}_{rt}$ using generalized natural growth operators.** In [5], the authors define an operator on $\mathcal{H}_{rt}$ that maps a forest to a single tree formed by connecting each root vertex of the forest to a new root vertex. Write this operator

$$B_+ : \mathcal{H}_{rt} \to \mathcal{H}_{rt} \ .$$

On a forest t_1t_2, this is defined

$B_+(t_1t_2) =$  .

Specifically, if $t_1 = t_2 = \bullet$, then

$B_+(\bullet, \bullet) =$  .

This can be extended by linearity to all of $\mathcal{H}_{rt}$.

Remark 1. The operator B_+ is a one Hochschild cocycle on $\mathcal{H}_{rt}$ [1].

The natural growth and the B_+ operator are related. For any tree t with root fertility n, there exists a set of trees $\{t_1 \dots t_n\}$ such that

$$t = B_+(t_1, \dots, t_n) \ .$$

Applying natural growth by t_0 gives

$$N_{t_0}(t) = B_+(t_0, t_1, \dots, t_n) + \sum_{i=1}^{n} B_+(t_1, \dots, N_{t_0}(t_i), \dots, t_n) \ . \tag{8}$$

In particular,

$$N_t(\bullet) = B_+(t) \tag{9}$$

Theorem 2. *The Hopf algebra of rooted trees, $\mathcal{H}_{rt}$, can be generated by elements formed by repeated application of operators of the form N_t to the tree with a single vertex.*

Proof. It is sufficient to show that any $t \in \mathcal{H}_{rt}$ can be written as a finite linear combination of elements in the set

$$\{N_{t_n}(\dots(N_{t_1}(\bullet))\dots) | n \in \mathbb{N}, t_i \text{ rooted tree}\} \ .$$

We proceed by induction on the fertility of the root vertex of t. Any rooted tree t with root fertility one can be written $t = B_+(t_1)$,

$$t = B_+(t_1) = N_{t_1}(\bullet) \ .$$

By induction, suppose all trees with root fertility $m-1$, that is of the form $B_+(t_1,\ldots,t_{m-1})$, can be written as a linear combination of trees formed by repeated application of general natural growth operators. By 8, the tree $t = B_+(t_1,\ldots,t_m)$ can be written

$$t = N_{t_m}(B_+(t_1,\ldots,t_{m-1})) - \sum_{i=1}^{m-1} B_+(t_1,\ldots,N_{t_m}(t_i),\ldots,t_{m-1}) \ .$$

Since every rooted tree t can be written as a linear combination of the set

$$\{N_{t_n}(\ldots(N_{t_1}(\bullet))\ldots)|n\in\mathbb{N}, t_i \in \mathcal{R}\} \ ,$$

it generates the Hopf algebra $\mathcal{H}_{rt}$. □

It is useful at this point to remark that the operator B_+ can be used to define the maps ϕ_t. Let $t = B_+(t_1\ldots t_n)$. Then

$$\phi_t(f) = \prod_{j=1}^{n} \phi_t(\phi^{i_j}(\bullet)) \prod_{j=1}^{n} \partial_{i_j} f \ . \tag{10}$$

Lemma 3. *Composing the maps ϕ_{N_t} and $\phi_{t'}$ gives*

$$\phi_{N_t}\phi_{t'} = \phi_{N_t(t')} \ .$$

Proof. Let $t' = B_+(t_1\ldots t_n)$. From equations 5 and 10,

$$\phi_{N_t}\phi_{t'}(h) = \phi^k(t)\partial_k\left(\prod_{j=1}^{n}\phi_t(\phi^{i_j}(\bullet))(\prod_{j=1}^{n}\partial_{i_j})\right)(h) =$$

$$\phi^k(t)\partial_k(\prod_{j=1}^{n}\phi_t(\phi^{i_j}(\bullet)))(\prod_{j=1}^{n}\partial_{i_j})(h) + \phi^k(t)(\prod_{j=1}^{n}\phi_t(\phi^{i_j}(\bullet)))(\partial_k\prod_{j=1}^{n}\partial_{i_j})(h) \ .$$

By equations 8 and 4, the second line evaluates to

$$\sum_{i=1}^{n}\phi_{B_+(t_1\ldots N_t(t_i)\ldots t_n)} + \phi_{B_+(tt_1\ldots t_n)} = \phi_{N_t(t')} \ .$$

□

2.3. **Sub Hopf algebras generated by natural growth.** One can generate sub Hopf algebras of $\mathcal{H}_{rt}$ by careful selection of a family of trees by which to grow.

Definition 10. Let S be a set of rooted trees. This defines a set of natural growth operators

$$\mathcal{N}(S) = \{N_t | t\in S\} \ .$$

Define $\mathcal{A}_S$ as the sub algebra of $\mathcal{H}_{rt}$ generated by the trees in S and the repeated application of operators in $\mathcal{N}(S)$. That is

$$\mathcal{A}_S = \mathbb{Q}[S][\{N_{t_n}(\ldots(N_{t_1}(s))\ldots)|n\in\mathbb{N}, t_i, s\in S\}] \ .$$

By construction, if $u \in \mathcal{A}_S$ and $t \in S$, then $N_t(u)\in\mathcal{A}_S$.

Theorem 3. *If $\mathbb{Q}[S] \in \mathcal{H}_{rt}$ is a Hopf algebra under Δ, then so is $\mathcal{A}_S$.*

Proof. It is sufficient to show that

$$\Delta : \mathbb{Q}[S] \to \mathbb{Q}[S]\otimes\mathbb{Q}[S] \quad \Rightarrow \quad \Delta : \mathcal{A}_S \to \mathcal{A}_S\otimes\mathcal{A}_S \ .$$

Since, $\mathbb{Q}[S]$ and $\mathcal{A}_S$ are graded and connected, this implies that they are Hopf algebras.

Consider $s, t\in S$, $N_t(s)\in\mathcal{A}_S$. By Theorem 1,

$$\Delta\circ N_t(s) = (N_t\otimes\mathbb{I})\circ\Delta s + \sum_{\substack{c_t \text{ admis. cut} \\ \text{of } t}} (P_{c_t}(t)\otimes N_{R_{c_t}(t)})\Delta s \ . \tag{11}$$

If $\mathbb{Q}[S]$ is a Hopf algebra, $\Delta(s) \in \mathbb{Q}[S] \otimes \mathbb{Q}[S]$. The first term in (1) is

$$(N_t \otimes \mathbb{I}) \circ \Delta s = \sum_{\substack{c_s \text{ admis. cut} \\ \text{of } s}} N_t(P_{c_s}(s)) \otimes R_{c_s}(s) \in \mathcal{A}_S \otimes \mathcal{A}_S.$$

The term $N_t(P_{c_s}(s)) \in \mathcal{A}_S$ since $P_{c_s}(s) \in \mathbb{Q}[S]$. Similarly, the second term in equation (11) can be written

$$\sum_{\substack{c_s \text{ admis. cut} \\ \text{of } s}} \sum_{\substack{c_t \text{ admis. cut} \\ \text{of } t}} P_{c_t}(t) P_{c_s}(s) \otimes N_{R_{c_t}(t)}(R_{c_s}(s)) \in \mathcal{A}_S \otimes \mathcal{A}_S \ .$$

Since the trees $t, s \in S$, the Hopf algebra structure of $\mathbb{Q}[S]$ implies that $R_{c_s}(s)$ and $R_{c_t}(t)$ are as well. Therefore $N_{R_{c_t}(t)}(R_{c_s}(s)) \in \mathcal{A}_S$.

For a set of trees $\{s, t_1 \dots t_n\}$ write $u_{s,t_1\dots t_n} \in \mathcal{A}_S$, defined

$$u_{s,t_1\dots t_n} = (N_{t_n}(\dots(N_{t_1}(s))\dots)) \ .$$

To calculate the coproduct of u

$$\Delta(u_{s,t_1\dots t_n}) = (N_{t_n} \otimes \mathbb{I}) \circ \Delta u_{s,t_1\dots t_{n-1}} + \sum_{\substack{c \text{ admis. cut} \\ \text{of } t_n}} (P_c(t) \otimes N_{R_c(t)}) \Delta(u_{s,t_1\dots t_{n-1}}) \ .$$

By induction, suppose that for all elements of the form $u_{s,t_1,\dots t_k}$ for $k < n$, and $s, t_i \in S$,

$$\Delta u_{s,t_1,\dots t_k} \in \mathcal{A}_S \otimes \mathcal{A}_S \ .$$

Then by similar arguments as above,

$$\Delta(u_{s,t_1\dots t_n}) \in \mathcal{A}_S \otimes \mathcal{A}_S \ .$$

Since the N_t are algebra homomorphisms, this extends to all $u \in \mathcal{A}_S$. □

Example 1. Let $S = \{\bullet\}$. The polynomial algebra $\mathbb{Q}[S]$ is a Hopf algebra since

$$\Delta\bullet = 1 \otimes \bullet + \bullet \otimes 1 \ .$$

The Hopf algebra generated by this set is $\mathcal{H}_{CK}$.

The rest of this section is devoted to computing the subHopf algebras defined by fan graphs.

Definition 11. Let F_i be the graph of weight i with $i-1$ leaves. These are called fan graphs.

For instance,



The function associated to each fan graph is

$$\phi^j(F_i) = \phi(\bullet)^{k_1} \cdots \phi(\bullet)^{k_{i-1}} (\partial_{k_1 \dots k_{i-1}} \phi(\bullet)^j) \ .$$

Let $S_k = \{F_i | i \leq k\}$. Using this notation, $\mathcal{H}_{CK} = \mathcal{A}_{S_1}$.

Theorem 4. *The algebras $\mathcal{A}_{S_k}$ are Hopf algebras.*

Proof. We need only check that $\mathbb{Q}[S_k]$ is a Hopf algebra. To see this, notice that for $n \leq k$,

$$\Delta F_n = 1 \otimes F_n + F_n \otimes 1 + \sum_{i=1}^{n-1} \binom{n-1}{i} \bullet^i \otimes F_{n-i-1} \in \mathbb{Q}[S_k] \otimes \mathbb{Q}[S_k] \ .$$

□

3. The Connes Moscovici Hopf algebras and their generalizations

In [6], the authors define a Hopf algebra $\mathcal{H}(n)$ on vector fields over F^+M, for M n-dimensional manifolds. This section recalls their construction, and generalizes it. For a clear explanation of Connes and Moscovici's construction of $\mathcal{H}(n)$, see [11].

For a n-dimensional smooth manifold, M, let

$$x^\mu : \mathbb{R}^n \to U \subset M$$

be a parametrization of a coordinate patch $U \subset M$. For $s \in \mathbb{R}^n$, write $y^\mu_i = \frac{\partial}{\partial s_i} x^\mu$. Let x^μ, y^μ_i form a basis of TU. In general, Greek indices correspond to the local coordinates on U and roman indices refer to the parametrizing indices in $\mathbb{R}^n$. Connes and Moscovici are only interested in orientation preserving diffeomorphisms on M, and thus only consider orientation preserving frames $F^+(M)$. In the one dimensional case, this means they can write

$$y(s) = e^{z(s)} \tag{12}$$

for some $z : \mathbb{R} \to M$. They define a connection $\mathfrak{gl}(n)$ valued one form on $TF^+(U)$,

$$\omega^l_j = (y^{-1})^l_\mu (dy^\mu_j + \Gamma^\mu_{\alpha,\beta} y^\alpha_j dx^\beta)$$

with Christoffel symbol $\Gamma^\mu_{\alpha,\beta}$. This connection defines a horizontal vector field in $TF^+(U)$

$$X_i = y^\mu_i (\partial_\mu - \Gamma^\beta_{\alpha,\mu} y^\alpha_j \partial^j_\beta) \ .$$

For $g \in C^\infty_c(U)$,

$$X_i(g) = \frac{\partial}{\partial s_i} g \ .$$

Further they define vector fields in $TF^+(U)$

$$Y^j_i = y^\mu_i \partial^j_\mu$$

which generate the $GL^+(n,\mathbb{R})$ action on $F^+(U)$.

In the one dimensional case, curvature, and thus Christoffel symbols, aren't defined. Instead define a function

$$\Gamma(x) = -\frac{1}{(\partial_s x)^2} \partial^2_s x \ .$$

If ψ is an orientation preserving local diffeomorphism on M with $\mathrm{Dom}(\psi) \subset U$, then

$$\Gamma(\psi) = -\frac{1}{(\partial_s \psi)^2} \partial^2_s \psi \ .$$

These function transform under coordinate change as Christoffel symbols do. Namely,

$$\Gamma(x)|_x = \partial_x \psi|_x \Gamma(\psi)|_{\psi(x)} + \frac{1}{\partial_x \psi} \partial^2_x \psi|_x \ . \tag{13}$$

In this case, the connection is defined

$$\omega = \frac{1}{y} dy + \Gamma(x) dx$$

with horizontal vector field

$$X = y\partial_x - y^2 \Gamma(x) \partial_y \ ,$$

and

$$Y = y\partial_y \ . \tag{14}$$

The vector fields X and Y act on the algebra generated by compactly supported functions on M crossed with the pseudo group of orientation preserving diffeomorphisms of M.

Definition 12. Let $\mathrm{Diff}^+(M)$ be the pseudo group of orientation preserving local diffeomorphisms on M. Define a group

$$\mathcal{A} = C_c^\infty(F^+(M)) \rtimes \mathrm{Diff}^+(M)$$

that is the semidirect product of compactly supported smooth functions on $F^+(M)$ with orientation preserving diffeomorphisms of M.

Any $\psi \in \mathrm{Diff}^+(M)$ lifts to a diffeomorphism of the frame bundle $\tilde{\psi} \in \mathrm{Diff}^+(F^+M)$

$$\tilde{\psi} : \{x^\mu, y_i^\mu\} \to \{\tilde{x}^\mu := \psi^\mu(x), \tilde{y}_i^\mu := y_i^\nu \partial_\nu \psi^\mu\} \ .$$

The group $\mathcal{A}$ is generated by the monomials

$$fU_\psi^* \in \mathcal{A} \quad f \in C_c^\infty(\mathrm{Dom}\tilde{\psi}) \quad \psi \in \mathrm{Diff}^+(U) \ .$$

The $*$ in the monomial corresponds to the contravariant multiplication

$$U_{\psi_1}^* U_{\psi_2}^* = U_{\psi_2\psi_1}^* \ .$$

Multiplication in the group is given by

$$(f_1 U_{\psi_1}^*)(f_2 U_{\psi_2}^*) = f_1 U_{\psi_1}^* f_2 U_{\psi_1}^{*-1} U_{\psi_1}^* U_{\psi_2}^* = f_1 \cdot (f_2 \circ \tilde{\psi}_1) U_{\psi_1\psi_2}^*$$

where $\cdot$ corresponds to pointwise multiplication. Writing $U_\psi^{*-1} = U_{\psi^{-1}}^*$, commutation with U_ψ^*

$$U_\psi^* f U_{\psi^{-1}}^* = f(p) \circ \tilde{\psi}(p) \tag{15}$$

results in changing the point of evaluation from $p \in F^+(U)$ to $\tilde{\psi}(p)$. The domain of $f_1 \cdot (f_2 \circ \tilde{\psi}_2)$ is

$$\mathrm{Dom}(\tilde{\psi}_1) \cap \tilde{\psi}_1^{-1}(\mathrm{Dom}\tilde{\psi}_2) \ .$$

The actions of X_i and Y on the monomials fU_ψ^* are given by

$$X_i(fU_\psi^*) = (X_i f)U_\psi^* \qquad Y(fU_\psi^*) = (Yf)U_\psi^* \ .$$

Before recalling their actions on products, we recall the push forwards of these vector fields. For a point $p \in F^+(U)$, and $\tilde{p} = \tilde{\psi}(p)$. Then

$$\tilde{\psi}_* Y_i^j|_{\tilde{p}} f = \tilde{y}_i^\mu \frac{\partial f(\tilde{p})}{\partial \tilde{y}_j^\mu} = y_i^\mu|_{\tilde{p}} \frac{\partial f(p)}{\partial y_j^\mu}|_{\tilde{p}} = Y_i^j|_{\tilde{p}} f \ . \tag{16}$$

That is, the vector field Y_i^j is invariant under orientation preserving diffeomorphisms. On the other hand,

$$\tilde{\psi}_* X_i|_{\tilde{p}} f = \frac{\partial}{\partial s_i}(f \circ \tilde{\psi}(p)) \neq \frac{\partial}{\partial s_i}|_{\tilde{p}}(f) \ .$$

Let $\tilde{\Gamma}_{\alpha,\beta}^\mu$ be the Christoffel symbol under the change of coordinates on M from x^μ to $\psi^\mu(x)$, Then

$$\tilde{\psi}_* X_i|_{\tilde{p}} = \tilde{y}_i^\mu \Big(\frac{\partial}{\partial \tilde{x}^\mu} - \tilde{\Gamma}_{\alpha,\beta}^\mu \tilde{y}_j^\alpha \frac{\partial}{\partial \tilde{y}_j^\beta}\Big) \ . \tag{17}$$

Any vector field V acting on a product of monomials $(f_1 U_{\psi_1}^*)(f_2 U_{\psi_1}^*)$, gives [11]

$$\begin{aligned} V|_p(f_1 U_{\psi_1}^*)(f_2 U_{\psi_1}^*) &= V|_p(f_1) \cdot (f_2 \circ \tilde{\psi}_1(p)) U_{\psi_2\psi_1}^* + f_1(p) \cdot \tilde{\psi}_{1*} V|_{\tilde{\psi}_1(p)} f_2 U_{\psi_2\psi_1}^* = \\ (V(f_1)U_{\psi_1}^*)(f_2 U_{\psi_2}^*) + (f_1 U_{\psi_1}^*) \cdot U_{\psi_1^{-1}}^* \tilde{\psi}_{1*} V|_{\tilde{\psi}_1(p)} f_2 U_{\psi_1}^* U_{\psi_2}^* &= (V(f_1)U_{\psi_1}^*)(f_2 U_{\psi_2}^*) + (f_1 U_{\psi_1}^*)(\tilde{\psi}_{1*} V|_p f_2 U_{\psi_2}^*) \ . \end{aligned} \tag{18}$$

By (16), since $\tilde{\psi}_{1*} Y_j^i|_p = Y_j^i|_p$,

$$Y|_p(f_1 U_{\psi_1}^*)(f_2 U_{\psi_2}^*) = (Y_j^i|_p(f_1 U_{\psi_1}^*)) f_2 U_{\psi_2}^* + (f_1 U_{\psi_1}^*)(Y_j^i|_p(f_2 U_{\psi_2}^*)) \ . \tag{19}$$

Applying (18) to X_i gives

$$X_i|_p(f_1 U_{\psi_1}^*)(f_2 U_{\psi_1}^*) = (X_i|_p(f_1)U_{\psi_1}^*)(f_2 U_{\psi_2}^*) + (f_1 U_{\psi_1}^*)((\tilde{\psi}_{1*} X_i)|_p f_2 U_{\psi_2}^*) \ .$$

Rewrite this

$$(X_i(f_1)U_{\psi_1}^*)(f_2 U_{\psi_2}^*) + (f_1 U_{\psi_1}^*)(X_i(f_2)U_{\psi_2}^*) + (f_1 U_{\psi_1}^*)((\tilde{\psi}_{1*} X_i - X_i)(f_2)U_{\psi_2}^*) \ . \tag{20}$$

By (17),

$$(\tilde{\psi}_{1*}X_i - X_i)|_p = \Gamma^{\mu}_{\alpha,\beta}|_p y^{\alpha}_j y^{\beta}_i (y^{-1})^k_{\mu} Y^j_k|_p - \tilde{\Gamma}^{\mu}_{\alpha,\beta}|_p (\tilde{y}^{\alpha}_j \tilde{y}^{\beta}_i (\tilde{y}^{-1})^k_{\mu} \tilde{Y}^j_k)|_p \ .$$

Definition 13. Write $\gamma^k_{ij}|_p(\psi_1)Y = (\tilde{\psi}_{1*}X_i - X_i)|_p$. Define a linear operator on $\mathcal{A}$, δ^k_{ij}, such that

$$\delta^k_{ij}(f_1 U^*_{\psi_1}) = (\gamma^k_{ij}|_p(\psi_1) f_1 U^*_{\psi_1}) \ .$$

For $a, b \in \mathcal{A}$, equation (20) gives

$$X_i(ab) = X_i(a)b + aX_i(b) + \delta^k_{ij}(a)Y^j_k(b) \ .$$

Connes and Moscovici show that

$$\delta^k_{ij}(ab) = \delta^k_{ij}(a)\delta^k_{ij}(b) \ .$$

When M is a one dimensional manifold, define $\delta_1 := \delta^1_{11}$, and $\gamma(\psi) = \gamma^1_{11}(\psi)$, which simplifies to

$$\gamma(\psi)|_p = y\Gamma(x)|_p - \tilde{y}\Gamma(\psi))|_p \ .$$

Writing $\psi' = \partial_x \psi$, and using equation (13), at the point $q = \tilde{\psi}(p)$,

$$\gamma(\psi)|_q = y\Gamma(x)|_q - y\Gamma(x)|_p + \frac{1}{\psi'}\frac{d\psi'}{ds}|_p \ . \tag{21}$$

The Hopf algebra $\mathcal{H}(n)$ is the universal enveloping algebra of the Lie algebra generated by Y^i_j, X_i, δ^i_{jk}, and their Lie products.

Remark 2. To simplify matters, Connes and Moscovici restrict to the domain where $\Gamma^i_{jk} = 0$, which ensures that

$$[X_i, X_j] = 0 \ .$$

If M is a one dimensional manifold, this is trivially true. We do not make such a requirement in the one dimensional case.

For the rest of this paper, we are only interested in the one dimensional case, where the Lie algebra of interest is generated by

$$Y, \quad X, \quad \{\delta_i | i \in \mathbb{N}\}$$

The Lie algebra has the commutation relations

$$[Y, X] = X; \quad [Y, \delta_i] = i\delta_i; \quad [X, \delta_i] = \delta_{i+1} \ .$$

The universal enveloping algebra of this Lie algebra is the Hopf algebra $\mathcal{H}(1)$. Commuting with the vector field X corresponds to the natural growth by a single vertex. The operators $\{\delta_i | i \in \mathbb{N}\}$ generate a Hopf algebra $\mathcal{H}_{CK}$ that is isomorphic to the sub Hopf algebra of rooted trees formed by applying the natural growth operator, $N_\bullet$, to the tree $\bullet$ from example 1.

3.1. The new Hopf algebra $\mathcal{H}_{rt}(1)$.

In this section, we enlarge $\mathcal{H}(1)$ to a Hopf algebra that incorporates vector fields X_t that correspond to natural growth by the tree t.

Since the vector field X in $\mathcal{H}(1)$ acts on δ_i as natural growth by $\bullet$, and $Y(\delta_1) = \delta_1$, we rename these maps to indicate the trees they correspond to in $\mathcal{H}_{rt}$. Specifically, define

$$X_\bullet := X \quad ; \quad \delta_\bullet := \delta_1 \ .$$

By this notation, define the function $\gamma(\psi) = \gamma_\bullet(\psi)$. Using the coordinates introduced in (12), and equation (14) we rewrite $X_\bullet$ in terms of the independent coordinates x, z. Specifically, $z = \log \frac{dx}{ds}$, $Y = \partial_z$, and

$$X_\bullet = e^z \partial_x - e^z \Gamma(x) \partial_z \ .$$

Notice that

$$\frac{dx}{ds} = e^z \quad ; \quad \frac{dz}{ds} = -e^z \Gamma(x) \ . \tag{22}$$

For the remainder of this paper, this defines the system of differential equations of interest. We apply the notation developed in section 2.1 to the equations in (22), with the index $i \in \{x, z\}$. In this context, rewrite

$$X_\bullet = \phi(\bullet)^i \partial_i \ .$$

The pushforward of $X_\bullet$ under an orientation preserving diffeomorphism $\tilde{\psi} \in \mathrm{Diff}^+(F^+M)$ gives the expression

$$\tilde{\psi}_* X_\bullet = \phi^x(\bullet)\partial_x + (\phi^z(\bullet) + \frac{d}{ds} \ \log \psi')\partial_z \ .$$

Similarly, for $q = \tilde{\psi}(p)$, $\gamma_\bullet(\psi)$ becomes

$$\gamma_\bullet(\psi)|_q = \phi^z(\bullet)|_p + (\frac{d}{ds} \log \psi')|_p - \phi^z(\bullet)|_q \ .$$

Definition 14. Define a function

$$\gamma_t(\psi) = \phi_t(\gamma_\bullet(\psi)) \ ,$$

according to defintion 4. This defines a linear operator δ_t on $\mathcal{A}$

$$\delta_t(fU_\psi) = \gamma_t(\psi) f U_\psi \ .$$

For a forest tt' of rooted trees, we write

$$\delta_{tt'} := \delta_t \delta_{t'} \ .$$

Lemma 4. *Consider the operator*

$$\phi_t : C_c^\infty(F^+M) \to C_c^\infty(F^+M) \ .$$

Write $t = B_+(t_1 \dots t_n)$. The pushforward of ϕ_t by $\tilde{\psi} \in \mathit{Diff}^+(F^+M)$ is given by

$$\tilde{\psi}_* \phi_t = \prod_{j=1}^n \phi_{t_j}(\tilde{\psi}_* \phi^{i_j}(\bullet)) \tilde{\psi}_* (\prod_{j=1}^n \partial_{i_j}) \ .$$

Proof. Let $\tilde{\psi} \in \mathrm{Diff}^+(F^+M)$. Recall that $\tilde{\psi} = (\psi, \log(\frac{d}{ds}\psi))$, for $\psi \in \mathrm{Diff}^+(M)$. Write $\tilde{\phi}$ as the map from $\mathcal{H}_{rt}$ to $C_c^\infty(F^+M)$ defined by coordinates defined by $\tilde{\psi}$. Define $\tilde{\phi}(t)$ accordingly. For $t = \bullet$, this is $\tilde{\psi}_* \phi^i(\bullet)$,

$$\tilde{\psi}_* \phi^i(\bullet) = \frac{d}{ds} \tilde{\psi}^i = \phi^j(\bullet) \partial_j \tilde{\psi}^i = \tilde{\phi}^i(\bullet) \ .$$

Since $\tilde{\phi}^i(\bullet) \frac{\partial}{\partial \tilde{\psi}^i} = \phi^i(\bullet)\partial_i$, we get

$$\tilde{\phi}^i(t) = \prod_{j=1}^n \phi^{i_j}(t_j) \prod_{j=1}^n \partial_{i_j} \tilde{\psi}_* \phi^i(\bullet)$$

for $t = B_+(t_1 \dots t_n)$. In other words

$$\tilde{\phi}^i(t) = \phi_t(\tilde{\psi}_* \phi^i(\bullet)) \ . \tag{23}$$

The pushforward of ϕ_t is

$$\tilde{\psi}_* \phi_t f = \phi_t(f \circ \tilde{\psi}) \ .$$

Applying this to equation (10) gives

$$\tilde{\psi}_* \phi_t = \prod_{j=1}^n \phi^{i_j}(t_j) \prod_{j=1}^n \partial_{i_j}(f \circ \tilde{\psi})$$

which can be rewritten using (23) as

$$\prod_{j=1}^n \tilde{\phi}(t_j) \prod_{j=1}^n \frac{\partial}{\partial \tilde{\psi}^{i_j}} f = \prod_{j=1}^n \phi_{t_j}(\tilde{\psi}_* \phi^i(\bullet)) (\prod_{j=1}^n \frac{\partial}{\partial \tilde{\psi}^{i_j}}) f \ .$$

□

The pushforward of ϕ_t leads to a useful expression for calculating its action for a product of monomials in $\mathcal{A}$.

Lemma 5. *The pushforward*

$$\tilde{\psi}_*\phi_t = \sum_{\substack{c \text{ admis. cut} \\ c \neq \text{ full}}} \gamma_{P_c(t)}(\psi)\phi_{R_c(t)}|_{\tilde{\psi}} \ .$$

For $R_c(t) = 1$, we write $\phi_1 = Y$.

Proof. Write $t = B_+(t_1 \dots t_n)$. By lemma 4 write

$$\tilde{\psi}_*\phi_t = \left(\prod_{j=1}^{k} \phi_{t_j}(\tilde{\psi}_*\phi^{i_j}(\bullet)) \prod_{j=1}^{k} \tilde{\psi}_*\partial_{i_j} \right) \ .$$

For subsets $I \subseteq \{1, \dots k\}$,

$$\tilde{\psi}_*\phi_t = \sum_I \left(\prod_{j \notin I} \phi_{t_j}(\tilde{\psi}_*\phi(\bullet) - \phi(\bullet)|_{\tilde{\psi}})^{i_j} \prod_{l \in I} \phi_{t_l}(\phi(\bullet)^{i_l} \circ \tilde{\psi}) \right) \prod_{j \notin I} (\tilde{\psi}_*\partial - \partial|_{\tilde{\psi}})_{i_j} \prod_{l \in I} \partial_{i_j}|_{\tilde{\psi}} \ . \tag{24}$$

Making the substitution

$$(\tilde{\psi}_*\phi(\bullet) - \phi(\bullet))^{i_j}(\tilde{\psi}_*\partial - \partial)_{i_j} = \gamma_\bullet(\psi)Y$$

into any of the summands of (24) corresponds to making an admissible cut of t with pruned forest $\prod_{j \notin I} t_j$. In the expression for $\tilde{\psi}_*\phi_t$, these appear as $\prod_{j \notin I} \phi_{t_j}(\gamma_\bullet(\psi))$. For a fixed set I, and a fixed $n \in I$, write the tree t_n in (24) $t_n = B_+(t_{n_1} \dots t_{n_r})$. For subsets $I' \subseteq \{1 \dots r\}$, use the definition of a pushforward and equation (24) to write the function

$$\phi_{t_n}(\phi^{i_n}(\bullet) \circ \tilde{\psi}) = \tilde{\psi}_*\phi_{t_n}(\phi^{i_n}(\bullet)) =$$

$$\sum_{I'} \left(\prod_{j \notin I'} \phi_{t_{n_j}}(\tilde{\psi}_*\phi(\bullet) - \phi(\bullet)|_{\tilde{\psi}})^{i_j} \prod_{l \in I'} \phi_{t_{n_l}}(\phi(\bullet)^{i_l} \circ \tilde{\psi}) \right) \prod_{j \notin I'} (\tilde{\psi}_*\partial - \partial|_{\tilde{\psi}})_{i_j} \prod_{l \in I'} \partial_{i_l}|_{\tilde{\psi}}(\phi^{i_n}(\bullet)) \ . \tag{25}$$

Substituting this expression into (24) corresponds to also taking an admissible cut of t_n. Therefore, the summands of $\tilde{\psi}_*\phi_t$, as given by equations (24) and (25) correspond to non-full admissible cuts of t. For each such cut, c, the function associated to the pruned forest by the map ϕ is of the form

$$\phi_{R_c(t)}\gamma_\bullet(\psi) = \gamma_{P_c(t)}(\psi) \ ,$$

an the root tree is

$$\phi_{R_c(t)}|_{\tilde{\psi}} \ .$$

□

We use this result to calculate the coproduct of δ_t.

Theorem 5. *The coproduct*

$$\Delta\delta_t = \sum_{c \text{ admis. cut}} \delta_{P_c(t)} \otimes \delta_{R_c(t)} \ .$$

Proof. By definition

$$\Delta\delta_t(fU_\psi)(gU_\eta) = \gamma_t(\eta \circ \psi) f \cdot (g \circ \psi) U_{\eta \circ \psi} \ ,$$

and

$$\gamma_t(\eta \circ \psi) = \phi_t(\gamma_\bullet(\eta \circ \psi)) \ .$$

From the coproduct of $\delta_\bullet$,

$$\gamma_\bullet(\eta \circ \psi) = \gamma_\bullet(\psi) + \gamma_\bullet(\eta) \circ \tilde{\psi} \ ,$$

so

$$\gamma_t(\eta \circ \psi) = \phi_t(\gamma_\bullet(\psi) + \gamma_\bullet(\eta) \circ \tilde{\psi}) = \gamma_t(\psi) + \tilde{\psi}_* \gamma_t(\eta) \ .$$

Rewriting $\tilde{\psi}_* \gamma_t(\eta) = \tilde{\psi}_* \phi_t(\gamma_\bullet(\eta))$, lemma 5 gives

$$\gamma_t(\eta \circ \psi) = \gamma_t(\psi) + \sum_{\substack{c \text{ admis. cut} \\ c \neq \text{ full}}} \gamma_{P_c(t)}(\psi) \gamma_{R_c(t)}(\eta) \circ \tilde{\psi} \ .$$

Therefore

$$\delta_t(fU_\psi)(gU_\eta) = \sum_{c \text{ admis. cut}} (\gamma_{P_c(t)}(\psi) fU_\psi)(\gamma_{R_c(t)}(\eta) gU_\eta) \ .$$

□

.

We use these linear operators to define a family of vector fields corresponding to natural growth.

Definition 15. Define a family of vector fields

$$X_t = \phi(t)^i \partial_i \ .$$

These act on $\mathcal{A}$ by $X_t(fU_\psi) = (X_t f)U_\psi$.

Remark 3. Notice that $X_t = \phi_{N_t}$ as defined in equation (5). By equation (9), one can write $X_t = \phi_{B_+(t)}$.

Theorem 6. *The coproduct of the vector fields X_t is the same as the coproduct of the natural growth operator on $\mathcal{H}_{rt}$. Namely,*

$$\Delta X_t = X_t \otimes 1 + \sum_{c \text{ admis. cut of } t} \delta_{P_c(t)} \otimes X_{R_c(t)} \ .$$

As with natural growth, we identify $Y = X_1$.

Proof. The coproduct of X_t is calculated by evaluating

$$X_t(fU_\psi)(gU_\eta) = (X_t f) \cdot (g \circ \psi) U_{\eta\psi} + f \cdot (\tilde{\psi}_* X_t g) U_{\eta\psi} \ .$$

Write $X_t = \phi_{B_+(t)}$, as in remark 3. Lemma 5 gives

$$\tilde{\psi}_* X_t = \sum_{\substack{c \text{ admis. cuts of } B_+(t) \\ c \neq \text{ full}}} \gamma_{P_c(B_+(t))}(\psi) \phi_{R_c(B_+(t))}|_{\tilde{\psi}} \ .$$

The edge set of $B_+(t)$, $E(B_+(t)) = E(t) \cup e$. A non-full admissible cut of $B_+(t)$ is either an admissible cut of t, or $c = e$. For any non-full admissible cut of $B_+(t)$, c, $R_c(B_+(t)) = B_+(R_c(t))$. If $c = e$, it is the full cut of t, and $R_c(t) = 1$. Therefore,

$$\psi_* X_t = \sum_{c \text{ admis. cuts}} \gamma_{P_c(t)} X_{R_c(t)}|_{\tilde{\psi}} \ .$$

□

This is of the same form as the coproduct of the natural growth operator N_t from Theorem 1.

Definition 16. Let $\mathcal{L}(1)$ be the Lie algebra spanned by the element Y, $\{X_t | t \text{ rooted tree}\}$ and $\{\delta_t | t \text{ rooted tree}\}$. Let $\mathcal{H}_{rt}(1) = \mathcal{U}(\mathcal{L}(1))$.

It remains to check the commutation relations on $\mathcal{L}(1)$.

Theorem 7. *The following commutation relations hold. Let t and t' be rooted trees.*

$$[Y, X_t] = |V(t)| X_t \quad ; \quad [X_t, \delta_{t'}] = \delta_{N_t(t')} \quad ; \quad [Y, \delta_t] = |V(t)| \delta_t$$
$$[X_t, X_{t'}] = X_{N_t(B_+(t'))} - X_{N_{t'}(B_+(t))} \quad ; \quad [\delta_t, \delta_{t'}] = 0 \ .$$

.

Proof. Recall that x and z are independent variables, and that $\Gamma(x)$ is only a function of x. Both $\phi^x(\bullet)$ and $\phi^z(\bullet)$ have a factor of e^z. Therefore, $Y(\phi^i(t)) = |V(t)|\phi^i(t)$. Writing

$$[Y, X_t](f) = Y(\phi^i(t))\partial_i(fU_\psi) = |V(t)|\phi^i(t)\partial_i(fU_\psi) = |V(t)|X_t(fU_\psi) \ .$$

Similarly,

$$[Y, \delta_t](fU_\psi) = |V(t)|\phi_t(\gamma_t(\psi)) = |V(t)|\delta_t(fU_\psi) \ .$$

To calculate

$$[X_t, \delta_{t'}](fU_\psi) = X_t(\phi_{t'}(\gamma_\bullet(\psi)))fU_\psi \ ,$$

use Lemma 3 to write

$$[X_t, \delta_{t'}](fU_\psi) = \phi_{N_t(t')}(\gamma_\bullet(\psi))(fU_\psi) = \delta_{N_t(t')}(fU_\psi) \ .$$

Similarly, the relation

$$[X_t, X_{t'}] = X_{N_t(B_+(t'))} - X_{N_{t'}(B_+(t))}$$

follows from lemma 3. Finally, $[\delta_t, \delta_{t'}] = 0$ follows from the fact that $\delta_t(fU_\psi) = \gamma_t(\psi)fU_\psi$. □

Finally, it is worth noting that the operators δ_t can be written in terms of a series of commutators of operators of the form $X_{t'}$ with $\delta_\bullet$.

Theorem 8. *The Lie algebra $\mathcal{L}(1)$ is generated by the set*

$$\{Y, \{X_t | t \in \mathcal{H}_{rt}\}, \delta_\bullet\}$$

and their commutators.

Proof. This is a corollary of theorem 2. Following the arguments presented there, write $t = B_+(t_1 \dots t_n)$. Assume by induction that for all t' with root fertility $< n$, $\delta_{t'}$ can be expressed in terms of linear combinations of commutators of $\delta_\bullet$ with a set $\{X_{t_i}\}$. Then

$$\delta_t(fU_\psi) = [X_{t_n}, \delta_{B_+(t_1 \dots t_{n-1})}](fU_\psi) - \sum_{i=1}^{n-1} \delta_{B_+(t_1 \dots N_{t_n}(t_i) \dots t_{n-1})}(fU_\psi) \ .$$

□

The crucial adjustment needed in order to generalize the Hopf algebra $\mathcal{H}(1)$ is that $\Gamma(x) \neq 0$. If $\Gamma(x) = 0$, then for any non-trivial rooted tree $t \neq \bullet$, $\phi^i(t) = 0$. Since $\phi^z(\bullet) = -e^z\Gamma(x)$, $\Gamma(x) = 0$ implies that the only non-zero term of the form $\partial_i\phi^j(\bullet)$ is $\partial_z\phi^x(\bullet)$. Therefore, $\Gamma(x) = 0$ implies that for any $t \neq \bullet$, only $\phi^x(t)$ is possibly non-zero. However, for $t = B_+(t_1 \dots t_n)$, one sees that

$$\phi^x(t) = \prod_{i=1}^{n} \phi^z(t_i)\partial_z^n(\phi^x(\bullet)) = 0.$$

Therefore, any generalization of this type for the algebras $\mathcal{H}(n)$, for $n \geq 1$ must incorporate the curvature of the base manifold M, as the commutator $[X_i, X_j] = R^k_{lij}Y^l_k$, for R^k_{lij} the curvature of M, and a torsion free connection ω^i_j on M. Towards this goal, work by [10] shows that planar rooted trees arise naturally in the case of flat connections with constant torsion.

References

[1] Christoph Bergbauer and Dirk Kreimer, *Hopf algebras in renormalization theory: Locality and Dyson-Schwinger equations from Hochschild cohomology*, IRMA Lectures in Mathematics and Theoretical Physics **10** (2006), 133 – 164, arXiv:hep-th/0506190v2.

[2] Christoper Brouder, *Runge-Kutta methods and renormalization*, European Physics Journal C **12** (2000), 521–534, arXiv:hep-th/9904014v1.

[3] John C. Butcher, *The numerical analysis of ordinary dierential equations*, Wiley-Blackwell, 1987.

[4] Arthur Cayley, *On the theory of the analytical forms called trees*, American Journal of Mathematics **4** (1881), 266–268.

[5] Alain Connes and Dirk Kreimer, *Hopf algebras, renormalization and non-commtative geometry*, Communications in Mathematical Physics **199** (1998), 203 – 242, arXiv:hep-th/9808042v1.

[6] Alain Connes and Henri Moscovici, *Hopf algebras, cyclic cohomology and the transverse index theory*, Communications in Mathematical Physics **198** (1998), 199–246, arXiv:math/9806109v1.

[7] Arne Dur, *Mobius functions, incidence algebras and power series representations*, Springer, 1986.

[8] Dirk Kreimer, *Dyson-Schwinger equations: From Hopf algebras to number theory*, In Universality and Renormalization (Ilia Binder and Dirk Kreimer, eds.), Field Institute Communications, vol. 50, American Mathematical society, 2006, arXiv:hep-th/0609004v2, pp. 225–248.

[9] Dirk Kreimer and Karen Yeats, *An Etude in non-linear Dyson-Schwinger Equations*, Nuclear Physics B. Proceedings Supplements **160** (2006), 116–121.

[10] A. Lunderwold and H. Munthe-Kaas, *On post-Lie algebra, Lie-Butcher series and moving frames*, arXiv:math-ph/1203.4738.

[11] Raimar Wulkenhaar, *On the ConnesMoscovici Hopf algebra associated to the diffeomorphisms of a manifold*, arXiv:math-ph/9904009v1.

UNIVERSITY OF HAMBURG, MATHEMATICS DEPARTMENT, BUNDESTRASSE, 55, 20146, HAMBURG, GERMANY

CALIFORNIA INSTITUTE OF TECHNOLOGY, 1200 E. CALIFORNIA BLVD., PASADENA, CA 91106, USA